

Whole Blood Potassium Polymorphism and Other Blood Electrolytes of Varamini Sheep in Iran

H. MORADI SHAHRBABAK¹, M. MORADI SHAHRBABAK AND H. MEHRABANI YEGANEH

Department of Animal Science, Faculty of Agriculture, University of Tehran, Karaj, Iran

¹Corresponding author's e-mail: hmoradis@ut.ac.ir

ABSTRACT

The whole blood potassium concentration has shown the bimodal distribution in sheep, which has been classified into LK and HK types; HK allele is recessive to LK with a single gene inheritance. This polymorphism showed different behavior in different environment, which could be due to adaptation process. This research was conducted on the Varamini breed research station, which has been located in a faculty of agriculture, university of Tehran. The whole Blood potassium concentration of 50 animals ranged from 6.5 - 28.06 m eq/L. The curve that the sheep could be divided into two subpopulation *Via* LK having 6.5 - 13.59 m eq/L of K⁺ and HK having 18.32 - 28.06 m eq/L of K⁺ with mean of 8.2 ± 0.25 m eq/L of K⁺ in the LK type and with mean of 23.54 ± 0.35 m eq/L of K⁺ in the HK types. The frequency of HK gene was found to be 0.91. Concentration of sodium, calcium and magnesium in whole blood were also determined. The relationship between potassium with sodium and calcium concentrations in whole blood of sheep was significant and negative estimated correlations were around -0.39 and -0.33, respectively which were significant. The mean of whole blood sodium and calcium concentration were 5147.4 and 86.54 ppm for LK and 4490 and 78.33 ppm for HK, respectively and all of differences were significant (p < 0.05). The mean of whole blood magnesium concentration was 31.73 ppm for LK and 35.12 ppm for HK, respectively and this differences was not significant (p < 0.05).

Key Words: Potassium; Whole blood; Polymorphism; LK; HK; Varamini sheep; Electrolytes; Na

INTRODUCTION

Evans and co-workers (Evans & King, 1955; Evans *et al.*, 1956; Evan & Mounib, 1957; Evans *et al.*, 1958; Evans, 1968) have shown that sheep of the same breed fall into two classes with respect to the K concentration within their red blood cells (R.B.C.S). The majority of individuals have low K (LK) cells and a minority has high K (HK) cells (Evans & Mounib, 1957), although the proportion of LK to HK animals may vary considerably between different breeds (Evans *et al.*, 1958). The authors presented convincing evidence for the claim that the cell type is genetically determined and that the genetic pattern can well be explained by they simply assumption of a single pair of alleles, the gene for LK being dominant over the gene for HK. several investigations have elucidated the physiological basis for the difference in R.B.C. K and (Na) content (Sheppard *et al.*, 1951; Joyce & Weatherall, 1958; Tosteson & Hoffman, 1960; Tosteson, 1963, 1966, 1967, 1969) the salient feature seems to be a high number of Na - K pump sites per unit cell membrane surface in HK cells and a low number in LK cells (Dunham & Hoffman, 1969; Hoffman, 1969; Lauf *et al.*, 1970). In addition, Tosteson and Hoffman (1960) found that the permeability of the membrane towards K and Na is different in the two cell types. The question as to whether there exist kinetic differences in the system for active cation transport in the two cell types is as Yet under discussion (Hoffman & Tosteson, 1969; Tosteson, 1969; Whittington & Blostein, 1971). In certain goat breeds a

similar dimorphism with respect to alkali cation content of red cells was found (Evans & Phillipson, 1957).

In cattle the alkali cation content of R.B.C.S is less well known. A study on Ayrshire cattle suggested that only on type of animals with LK cells (with a mean value of 20 m-mole/L. cells & a range from 12 to 39 m-mole/L. cells) can be found (Evans & Phillipson, 1957), although the contention exists that a bimodal distribution is universally encountered in ruminants (Evans, 1968). We therefore decided to study the problem using Swiss cattle breeds. The survey included mainly animals of the Simmenthal breed but also a few Swiss brown (braunvieh), Swiss black spotted (Freiburger Schwarzfleckvieh) and others. Since analysis of the results revealed no difference between breeds, the figures from all animals tested were pooled. An attempt was made to describe quantitatively the differences in passive permeability, Na - K pump activity and Na + K activated membrane ATP ase by measuring net cation fluxes in cold, net cation fluxes at 37 C and phosphate liberation from ATP by isolated membranes. This paper describes the genetic polymorphism of potassium level and its relationship with other blood electrolytes in Varamini Sheep in Iran.

MATERIALS AND METHODS

This study conducted in the Varamini Research station that located in the shahrbabak -Iran. The material for the present study consisted of 50 sheep. Since it is known that LK sheep and cattle are born With High R.B.C. K-

concentration, which falls to a low steady level during the first weeks of life (Caldwell *et al.*, 1947; Widdas, 1954; Wise Wright *et al.*, 1958; Tosteson & Moulton 1959; Evans & Blunt, 1961; Blencher 1961; Tosteson, 1966) only animal above 4 months of age were used. Climate conditions and feeding habits were approximately similar for all animals tested.

Sheep were selected randomly and blood was collected into heparinized vials as anticoagulant and from the jugular vein and brought to the laboratory in an ice cooled isolating box.

Determination of potassium and blood electrolytes (Na, Ca & Mg) was carried out by using a spectrophotometer of atomic absorption system (shimadzu). Evans *et al.* (1956) and Khattab *et al.* (1964) have reported the plasma concentration to be same in both LK and HK sheep and the differences in whole blood potassium was attributed to the difference in red blood cell K⁺ concentration. Hence only the Whole blood K⁺ concentration was estimated in the present study and this was taken as reflecting the K⁺ concentration in the erythrocytes.

Since potassium is the cation of Intracellular and there is in the erythrocytes, the erythrocyte must be lysed for measurement of potassium. For K determination whole blood were diluted with water distiller and we diluted of 0.1 mm of sample in ratio 1:500 and then potassium concentration in the whole blood was determined in the spectrophotometer of atomic absorption system (shimadzu).

For Na determination whole blood were diluted with water distiller and we diluted of sample in ratio 1:4500 and then sodium concentration in the whole blood was determined in the spectrophotometer of atomic absorption system (shimadzu).

For Mg (magnesium) and Ca (calcium) determination whole blood were diluted with water distiller and it's diluted of sample in ratio 1:30 and then sodium concentration in the whole blood was determined in the spectrophotometer of atomic absorption system (shimadzu).

Concentration of the electrolytes in the whole blood was determined by the spectrophotometer of atomic absorption system (shimadzu) in ppm and then unit concentration (ppm) of potassium convert to meq/L for determination of polymorphism.

Frequency of gene and genotype of whole blood potassium polymorphism were calculated by collected data and then LK and HK samples were determined by curve normal distribution (bimodal distribution has showed two phenotype in whole blood potassium and we can separate these phenotypes) (Suzuki, A. *et al.*, 1991).

Analysis variance and correlation of blood potassium polymorphism with other electrolytes were calculated by GLM in SAS program, respectively.

Phenotypic correlations between potassium concentration and other electrolytes were calculated by Pearson method in correlation procedure in SAS program.

RESULTS

The whole Blood potassium concentration of 50 animals ranged from 6.5 - 28.06 m eq/L. The frequency distribution of blood potassium concentration in the sheep population is presented as a frequency curve. It is seen from the bimodal nature. The curve that the sheep could be divided into two sub-population *Via* LK having 6.5 - 13.59 m eq/L of K⁺ and HK having 18.32 - 28.06 m eq/L of K⁺ with mean of 8.2 ± 0.25 m eq/L of K⁺ in the LK type and with mean of 23.54 ± 0.35 m eq/L of K⁺ in the HK types. The bimodal nature seen in the study is in agreement with that observed by Evans (1954), Evans and Mounib (1957), Pater and Suska (1962), Meyer (1963), Khattab *et al.* (1963).

The frequency distribution of LK and HK phenotypes and their gene frequency obtained in the population are given in Table I. It seen that 82% of Varamini sheep are of HK type while 18% is of LK type. The frequency of HK gene was found to be 0.91 (Table I). Concentration of sodium, calcium and magnesium in whole blood were also determined, the mean of whole blood sodium and calcium concentration were 5147.4 and 86.54 ppm for LK and 4490 and 78.33 ppm for HK, respectively and all of differences were significant ($p < 0.05$). The mean of whole blood magnesium concentration was 31.73 ppm for LK and 35.12 ppm for HK, respectively and this differences was not significant ($p < 0.05$) (Table II & III).

Remarkable differences significant in whole blood of calcium and magnesium concentrations were recognized between LK and HK types ($P < 0.05$) (Table III).

The relationship between potassium with sodium and calcium concentrations in whole blood of sheep was significant and negative estimated correlations were around - 0.39 and -0.33, respectively which were significant (Table IV).

DISCUSSION

Although the HK allele is recessive allele but frequency of HK allele is 0.9. In the literature showed that animals with HK phenotype have active Na-k-pump (i.e. concentration of Na & k in cell regulate with consume energy) but the animal with LK phenotype, concentration of Na and K regulate with simple diffusion and Na-K-pump is semi active. The relationship between potassium polymorphism and whole blood sodium concentration is negative. The animal with LK type have a high level of sodium but the animal with HK type Have a low level of sodium and this result return to performance of NA-K-pump activity that discussed above. Therefore to regulate of Na K concentration in cell, the animal with HK genotype can better survive in this environment.

Acknowledgement. We thank the University of Tehran. We are also thankful to Organization of Jihade- keshavarzi kerman and sharbabak the Miss Eng. Khoshvaghti for analysis of blood samples.

Table I. Determination gene and genotype of blood potassium polymorphism

Phenotype	Number of observation	% observation	Gene frequency
HK	41	82	(HK) 0.91
LK	9	18	
Total	50	100	

Table II. Comparison of Duncan's mean of Na within potassium types of Varamini sheep

Phenotype	number	Means (ppm)
LK	9	5147.4 ^{a*}
HK	41	4490 ^b

* The different letter show the significant different t (P<0.01)

Table III. Comparison of Duncan's mean of Ca (calcium) and Mg (magnesium) within potassium types of Varamini sheep

Phenotype	number	Ca Means (ppm)	Mg means (ppm)
LK	9	86.54 ^{b*}	31.73 ^b
HK	41	78.33 ^a	35.12 ^a

* The same letter show the non significant different t (P<0.05)

Table IV. Phenotypic correlations between potassium concentration and other electrolytes

Electrolytes	Ca (ppm)	Na (ppm)	Mg (ppm)
correlation coefficient	-0.33	-0.39	-0.28
level of significant	0.03	0.01	0.08
number of sheep	50	50	50

REFERENCES

Campbell, E.H., J.C. Ellory and J.C. Gibson, 1998. Anti-L antibody and the regulatory phosphorylation cascade of Lk sheep red blood cell. *J. Physiol.*, 27b: 357-64

Christinaz, P. and H.J. Schatzmann, 1972. High potassium and low potassium erythrocytes in cattle. *J. Physiol.*, 224: 391-406

Dunham, P.B., 1976a. Two populations of antibodies affecting cation transport in LK erythrocytes of sheep and goats. *Bioch. Et. Biophys. Acta*, 443: 219-26

Dunham, P.B. and R. Blostein, 1976b. Active potassium transport in reticulocytes of High-K⁺ and Low-K⁺ sheep. *Bioch. Et. Biophys. Acta*, 455: 749-58

Dunham, P.B. and R. Blostein, 1997. L-antigens of sheep red blood cell membranes and modulation of ion transport. *American J. Physiol.*, 272: 357-68

Dunham, P.B. and J.C. Ellory, 1981. Passive potassium transport in low potassium sheep red cells: Dependence upon cell Volume and Chloride, 318: 511-30

Ellory, J.C. and E.M. Tucker, 1970. Active potassium transport and the L and M antigens of sheep and goat Red cells. *Bioch. Et. Biophys. ACTA.*, 58: 160-68

Evans, J.V., 1954. Electrolyte concentration in red blood cells of British breeds of sheep. *Nature*, 174: 931-2

Evans, J.V., 1957. Water metabolism in the sheep. *Nature*, 180: 931

Evans, J.V., H. Harris and F.L. Warren, 1958. Hemoglobin and potassium blood types in some Non-British Breeds of sheep and in certain rare British Breeds. *Nature*, 182: 320-7

Evans, J.V., J.W.B. King, B.L. Cohen, H. Harris and F.L. Warren, 1956. Genetic of Hemoglobin and blood potassium differences in sheep. *Nature*, 849-50

Evans, J.V. and J.W. King, 1955. Genetic control of sodium and potassium concentration in the red blood cell of sheep. *Nature*, 176: 171-2

Khan, M.S. and G.C. Taneja, 1983. Lk gene and wool production characteristics in rajasthan desert sheep. *Indian. J. Anim. Sci.*, 53: 898-901

Khan, M.S., 1997. *Biochemical Model for Sustained Animal Production in Arid Ecosystem*. Pp: 353-62. Central Arid Zone Research Institute, India

Krishnamurthy, U.S. and V. Rathnasabapathy, 1977. Genetics of blood potassium in Nilagiri merino and their cross-bred sheep. *Indian Vet. J.*, 54: 36-41

Lipecka, C., M. Pieta and T. Gruszecki, 1991. Genetic polymorphism of blood potassium in various breeds and strains of sheep. *Genetica Polonica*, 32: 113-7

Lipecka, C., M. Pieta and T. Gruszecki, 1994. The potassium level in the blood of sheep and their productivity. *J. Anim. Feed. Sci.*, 3: 89-96

Moradi Shahrabak, H.M., M. Shahrabak, G. Rahimi and A. Nikkhah, 2002. Estimation of body weight traits (co) variance components and genetic parameters in kermani sheep and study of blood potassium polymorphism and its relationship with production trait. *M.Sc. Thesis*, Department of Animal Science, University of Tehran.Karaj, Iran

Pittman, S.J., J.C. Ellory, E.M. Tucker and C.I. Newbold, 1990. Identification of a 25 Kda polypeptide associated with the L-antigen in low potassium-type sheep red cells. *Bioch. Et. Biophys. Acta*, 1022: 408-10

Reddy, V.R.C. and A.R. Krishnan, 1988. Blood potassium types and their relationship with body weight in sheep. *Cheiron*, 17: 114-20

Reddy, V.R., V.P. Padmanabha and K.K. Reddy, 1990. Distribution of Blood potassium types and their relationship with Growth traits in Nellore and Nellore X Dorset Half-Bred sheep. *Indian Vet. J.*, 67: 420-4

Singh, L.B., P.K. Dwarakanath, A.V. Rakim and S.P. Prasad, 1979. Blood electrolyte (K & Na) and Hb-Variants in karakul sheep. *Indian Vet. J.*, 56: 554-6

Suzuki, A. and K. Sugawara, 1991. Frequency of appearance of low and high potassium types and mineral concentration in whole blood of sheep. *Tohoku J. Agric. Res.*, 41: 57-60

Swenson, M.J. and W.O. Reece, 1996. *Duckes Physiolog*, Pp: 24-7

Whittington, E.S. and R. Blostein, 1971. Comparative properties of high potassium and low potassium sheep erythrocyte membrane sodium activated adenosine triphosphatase. *J. Biol. Chem.*, 246: 3518-23

Wright, R.D., T.R. Bradley, J.F. Nelson and J.P. Coghlan, 1985. Changes in the potassium concentration and metabolism of red blood cells of the lamb. *Nature*, 182: 1742-3

Wright, R.D., T.R. Bradley, J.F. Nelson and J.P. Coghlan, 1985. Changes in the potassium concentration and metabolism of red blood cells of the lamb. *Nature*, 182: 1742-3

(Received 13 February 2006; Accepted 20 June 2006)