

# Induced Mutations for Yield Contributing Traits in Green Gram

SAMIULLAH KHAN<sup>1</sup> AND MOHD. RAFIQ WANI

*Mutation Breeding Laboratory, Department of Botany, Aligarh Muslim University, Aligarh – 202 002 (U.P.), India*

<sup>1</sup>Corresponding author's e-mail: khansami2004@rediffmail.com

## ABSTRACT

In the present investigation, variability induced by ethylmethane sulphonate (EMS) for four quantitative traits viz., fertile branches/plant, pods/plant, 100-seed weight (g) and yield/plant (g) was studied in M<sub>2</sub> and M<sub>3</sub> generations of green gram (*Vigna radiata* L. Wilczek) var. Asha. A significant shift in mean values in the positive direction was observed for all the quantitative traits studied. High values of heritability and genetic advance over the control indicate the possibility of inducing desirable mutations for polygenic traits accompanied with effective selection in M<sub>2</sub> and later generations. Yield/plant was positively correlated with various yield contributing traits, but did not show any association with seed protein content.

**Key Words:** Green gram; EMS; Quantitative traits; Correlations; Protein content

## INTRODUCTION

Variability in base population becomes essential when breeding objectives are more complex. Main interests of a plant breeder are quantitative traits, which are controlled by polygenic interactions. A series of experiments carried out with various crops have established that chemical mutagens induce polygenic variability (Kharkwal, 1999; Rajput *et al.*, 2001; Singh & Singh, 2001; Khan *et al.*, 2004). Since green gram is a self-pollinating crop, limited variability is available for its improvement programmes. Induced mutagenesis thus seems to be an ideal methodology for the induction of desirable genetic variability.

The present study was undertaken to evaluate the extent of genetic variability in M<sub>2</sub> and M<sub>3</sub> generations for the four quantitative traits of green gram var. Asha following mutagenesis with ethylmethane sulphonate (EMS). An attempt was also made to find out the association between yield and yield components.

## MATERIALS AND METHODS

A field experiment was conducted during kharif season of 2000/01, 2001/02 and 2002/03 at the Agricultural Farm, Aligarh Muslim University, Aligarh, India. Uniform and healthy seeds of green gram (*Vigna radiata* L. Wilczek) var. Asha were presoaked in distilled water for 9 h followed by the treatment with EMS (0.1 – 0.3%), prepared in phosphate buffer of pH 7 for 6 h. Seeds soaked in distilled water were used as control. After completion of treatment period for 6 h, the treated seeds were washed thoroughly in running tap water to reduce the residual effect of the mutagen sticking to the seed coat. The treated and control seeds were sown in the field in complete randomized block design (CRBD). Seeds harvested from individual M<sub>1</sub> plants were sown as M<sub>2</sub> families in three replications in the field. For raising M<sub>3</sub> generation, such 10 M<sub>2</sub> progenies were

selected, which showed significant deviations in mean values in the positive direction from the mean values of control, particularly for the yield and yield components in M<sub>2</sub> generation. Seeds from each selected M<sub>2</sub> progeny were bulked by taking an equal amount of seeds from all M<sub>2</sub> plants from a single M<sub>2</sub> progeny and thoroughly mixed. A random sample of this bulk was sown to obtain M<sub>3</sub> progeny.

ANOVA was performed to test the significance of variance. Heritability ( $h^2$ ) in broad-sense and phenotypic correlation among various character pairs were calculated by the formulae suggested by Singh and Chaudhary (1985) and Khan and Khanum (1994), respectively and the genetic advance by the method outlined by Johnson *et al.* (1955). The protein content of seeds was determined by the method of Lowry *et al.* (1951).

## RESULTS AND DISCUSSION

The data recorded for various quantitative traits in green gram are presented in Tables I and II. It is clear from the tables that the mean values increased significantly for all the yield contributing traits under study. Higher values of mean for number of fertile branches, number of pods/plant and total plant yield were recorded with 0.2% EMS treatment in both M<sub>2</sub> and M<sub>3</sub> generations. It has been observed that during mutagenesis, if mutations occur at random for the quantitative traits, no significant change is expected in the mean values. However, the shift in mean values in the positive direction indicates that more positive mutations have occurred for these traits. Plant yield was found to be associated with the increase in the number of fertile branches, pods, and 100-seed weight. A high degree of correlation between number of pods/plant and number of fertile branches was reported earlier by Khan and Siddiqui (1997) in mungbean and Kharkwal (2003) in chickpea. The values for genotypic variability, heritability and genetic advance also showed an increase over control in most of the

**Table I. Estimates of mean values and genetic parameters for different yield contributing traits in M<sub>2</sub> generation of green gram var. Asha**

Treatment	Mean±S.E.	Shift in $\bar{X}$	GCV (%)	h <sup>2</sup> (%)	GA(% of $\bar{X}$ )
<b>Fertile branches / plant</b>					
Control	6.77±0.18	-	7.30	20.35	6.95
0.1% EMS	7.34±0.21	+0.57*	10.81	25.90	13.89
0.2 % EMS	7.83±0.27	+1.06*	18.67	54.08	36.25
0.3% EMS	7.10±0.25	+0.33*	11.80	25.00	11.43
<b>Pods / plant</b>					
Control	47.97±1.60	-	1.38	15.06	1.53
0. 1% EMS	49.61±0.25	+1.64*	11.10	60.55	16.15
0. 2% EMS	51.70±0.26	+3.73*	3.40	69.10	22.50
0.3% EMS	48.33±0.25	+0.36*	7.50	66.85	7.62
<b>100 seed-weight (g)</b>					
Control	4.10±0.06	-	3.40	14.30	2.40
0.1% EMS	4.55±0.03	+0.45*	7.10	33.36	10.55
0.2% EMS	4.20±0.05	+0.10	6.15	42.75	10.38
0.3% EMS	4.29±0.06	+0.19	5.21	21.75	5.85
<b>Yield / plant (g)</b>					
Control	8.60±0.19	-	4.36	44.80	4.80
0.1% EMS	9.02±0.21	+0.42*	7.35	60.00	14.75
0.2% EMS	9.50±0.23	+0.90*	17.41	48.70	32.07
0.3% EMS	8.87±0.19	+0.27*	9.10	45.98	16.20

\* = Significant at p = 0.05; GCV = Genotypic coefficient of variation; h<sup>2</sup> = Heritability, GA = Genetic advance.

**Table II. Estimates of mean values and genetic parameters for different yield contributing traits of the mutants isolated in M<sub>3</sub> generation of green gram var. Asha**

Treatment	Mean±S.E.	Shift in $\bar{X}$	GCV (%)	h <sup>2</sup> (%)	GA(% of $\bar{X}$ )
<b>Fertile branches / plant</b>					
Control	6.80±0.17	-	6.06	17.53	6.66
0.1% EMS	7.56±0.21	+0.76*	20.49	72.07	45.76
0.2 % EMS	7.96±0.22	+1.16*	21.86	76.71	50.36
0.3% EMS	7.44±0.19	+0.64*	17.01	66.11	36.30
<b>Pods / plant</b>					
Control	47.94±0.15	-	2.56	14.24	2.50
0. 1% EMS	52.13±0.73	+4.19*	12.46	84.55	30.25
0. 2% EMS	53.23±0.72	+5.29*	11.86	83.14	28.48
0.3% EMS	47.73±0.57	-0.21	8.15	68.14	17.71
<b>100 seed-weight (g)</b>					
Control	4.03±0.10	-	5.54	14.28	5.41
0.1% EMS	4.84±0.17	+0.81*	7.73	14.89	7.83
0.2% EMS	4.73±0.15	+0.70*	8.72	19.10	9.93
0.3% EMS	3.90±0.13	-0.13	9.59	22.58	11.88
<b>Yield / plant (g)</b>					
Control	8.53±0.02	-	6.93	44.87	12.20
0.1% EMS	9.13±0.15	+0.60*	12.59	73.74	28.47
0.2% EMS	9.86±0.33	+1.33*	21.49	63.59	45.03
0.3% EMS	8.63±0.24	+0.10	17.30	60.92	35.58

\* = Significant at p = 0.05; GCV = Genotypic coefficient of variation; h<sup>2</sup> = Heritability, GA = Genetic advance.

treatments. The high heritability estimates in the quantitative characters have been found to be useful for selecting suitable types based on their phenotypic performance. The data, in general, indicate a relatively higher estimated heritability (broad-sense) for various quantitative traits in M<sub>3</sub> generation. The increased heritability values in M<sub>3</sub>, in comparison to M<sub>2</sub> generation, may be due an increased homozygosity of the genes involved. Johnson *et al.* (1955) advocated that heritability estimates along with genetic advance is usually more helpful than the heritability value

alone in predicting the resultant effects of selection. Genetic advance is indicative of the expected genetic progress for a particular trait under suitable selection procedure (Kaul & Garg, 1982) and consequently carries much significance in self-pollinated crops. The results are in conformity with other reports (Khan *et al.*, 2001; Singh *et al.*, 2001; Raut *et al.*, 2004). In the present study, the higher values of heritability and genetic advance suggest that the variability so evolved can be effectively exploited for further genetic improvement of green gram.

**Table III. Phenotypic correlation between various character pairs of the mutants isolated in M<sub>3</sub> generation of green gram**

Treatment	Fertile branches/plant vs. pods/plant	Fertile branches/plant vs. yield/plant	Pods/plant vs. yield/plant	100-seed weight vs. yield/plant
Control	+ 0.022	+ 0.161	+ 0.330	- 0.301
0.1% EMS	+ 0.482**	+ 0.326	+ 0.449*	- 0.205
0.2% EMS	+ 0.522**	+0.487**	+ 0.292	- 0.550
0.3% EMS	+ 0.045	+ 0.186	+ 0.392*	- 0.102

\*,\*\*Significant at 5% and 1% levels of probability, respectively.

**Table IV. Total seed protein content (%) of the mutants isolated in M<sub>3</sub> generation of green gram**

Treatment	Mean ±S.E.	Seed protein content (%)		CV (%)	Yield/plant vs. Seed protein content (r)
		Shift in $\bar{X}$	Range		
Control	21.38±0.33	-	20.10-22.10	3.50	- 0.183
0.1% EMS	22.92±0.85	+ 1.54	19.80-24.85	8.28	- 0.723
0.2% EMS	23.18±0.97	+ 1.80	19.55-24.95	9.40	- 0.059
0.3% EMS	23.02±0.74	+ 1.64	20.08-23.40	7.17	- 0.674

CV = Coefficient of variation; r = Correlation coefficient

Correlation coefficients among four characters under study were estimated at phenotypic level (Table III). In all the mutagenic concentrations, the correlations were found to be higher than the control. Yield/plant was found to have significant positive correlation with number of pods/plant and fertile branches/plant. These findings are in conformity with those reported by Vijayalakshmi *et al.* (2000) in chickpea. 100-seed weight was not correlated with yield per plant in the present study. However, Amaranath *et al.* (1990) and Singh and Singh (1999) reported positive correlation of 100-seed weight with total plant yield in soybean.

Seed protein content of the mutants did not alter significantly in any of the EMS treatments studied (Table IV). However, the coefficient of variation was higher in mutagenic treatments as compared to control, indicating that improvement in seed protein content is possible. Seed protein did not show any association with yield/plant, indicating the independent genetic control of protein content and total plant yield. Sandhu *et al.* (1979) suggested that the lack of association between grain yield and grain protein can be easily combined in a single genotype.

**Acknowledgements.** We are grateful to the Chairman, Department of Botany, Aligarh Muslim University, Aligarh for providing necessary research facilities.

## REFERENCES

Amaranath, K.C.N., S.R. Viswanatha and S.R. Chennakeshava, 1990. Phenotypic and genotypic correlation coefficient of some quantitative characters in soybean (*Glycine max* L. Merrill). *Mysore J. Agric. Sci.*, 24: 445-9

Johnson, H.W., H.F. Robinson and R.E. Comstock, 1955. Estimates of genetic and environmental variability in soybeans. *Agron. J.*, 47: 314-8

Kaul, M.L.H. and R. Garg, 1982. Radiation genetic studies in garden pea. I. Genetic variability, inter-relationships and path analysis in protein rich genotypes. *Biol. Zbl.*, 101: 271-82

Khan, I.A. and A. Khanum, 1994. *Fundamentals of Biostatistics*. Ukaaz Publishers, Hyderabad, India

Khan, S. and B.A. Siddiqui, 1997. Genetic variability and correlation studies in yield and yield components in mungbean (*Vigna radiata* L. Wilczek). *J. Indian Bot. Soc.*, 71: 69-71

Khan, S., M.R. Wani and K. Parveen, 2004. Induced genetic variability for quantitative traits in *Vigna radiata* (L.) Wilczek. *Pakistan J. Bot.*, 36: 845-50

Khan, S., M. Rehman, M. Bhat and B.A. Siddiqui, 2001. MMS induced biological damage and polygenic variability in green gram (*Vigna radiata* L. Wilczek). *Legume Res.*, 23: 126-9

Kharkwal, M.C., 1999. Induced mutations in chickpea (*Cicer arietinum* L.) III. Frequency and spectrum of viable mutations. *Indian J. Genet.*, 59: 451-64

Kharkwal, M.C., 2003. Induced mutations in chickpea (*Cicer arietinum* L.) VI. Significance of induced altered correlations. *Indian J. Genet.*, 63: 219-24

Lowry, O.H., N.J. Rosebrough, A.L. Farr and R.J. Randall, 1951. Protein measurement with folin phenol reagent. *J. Biol. Chem.*, 193: 265

Rajput, M.A., G. Sarwar and K.A. Siddiqui, 2001. Development of high yielding mutants in lentil. *Mut. Breed. Newslet.*, 45: 35-6

Raut, V.K., J.V. Patil and V.L. Gawande, 2004. Correlation and path analysis for quantitative traits in chickpea. *Indian J. Pulses Res.*, 17: 82-3

Sandhu, T.S., B.S. Bhullar, H.S. Cheema and A.S. Gill, 1979. Variability and inter relationships among grain protein, yield and yield components in mungbean. *Indian J. Genet.*, 39: 480-4

Singh, G., P.K. Sareen, R.P. Saharan and A. Singh, 2001. Induced variability in mungbean (*Vigna radiata* L. Wilczek). *Indian J. Genet.*, 61: 281-2

Singh, J. and B. Singh, 1999. Genetic variability and correlation studies in soybean (*Glycine max* L. Merrill). *J. Oil Seeds Res.*, 16: 118-20

Singh, M. and V.P. Singh, 2001. Genetic analysis of certain mutant lines of urd bean for yield and quality traits in M<sub>4</sub> generation. *Indian J. Pulses Res.*, 14: 60-2

Singh, R.K. and B.D. Chaudhary, 1985. *Biometrical Methods in Quantitative Genetic Analysis*. Kalyani Publishers, Ludhiana, India

Vijayalakshmi, N.V.S., K. Jagdish and T.N. Rao, 2000. Variability and correlation studies in Desi, Kabuli and intermediate chickpeas. *Legume Res.*, 23: 232-6

(Received 15 December 2005; Accepted 20 May 2006)