



**Full Length Article**

# Effectiveness of *Chrysoperla carnea* (Stephens) (Neuroptera: Chrysopidae) on the Population of *Bemisia tabaci* (Homoptera: Aleyrodidae) in Different Cotton Genotypes

KHURAM ZIA, FAISAL HAFEEZ<sup>1</sup>, RASHID RASOOL KHAN, MUHAMMAD ARSHAD AND UNSAR NAEEM-ULLAH

Department of Agri. Entomology, University of Agriculture, Faisalabad, Pakistan

<sup>1</sup>Corresponding author's e-mail: faisalhafeez143@yahoo.com

## ABSTRACT

Present study was conducted to determine the effectiveness of *Chrysoperla carnea* in sustainable biological control programs. The experiment was conducted on six advanced genotypes of cotton cultivars viz. MNH-552, CIM-707, SLH-284, IRFH-901, CIM-496 and FH-115. *Bemisia tabaci* (Genn.) (Homoptera: Aleyrodidae) was used as a prey in the presence and absence of its natural predator, *Chrysoperla carnea* (Stephens) (Neuroptera: Chrysopidae). *Chrysoperla carnea* depicted maximum reduction of whitefly population (57.35%) in 4<sup>th</sup> week of August in MNH-552 followed by 2<sup>nd</sup> week of September in MNH-552 (57.14%), 3<sup>rd</sup> week of August in FH-115 (55.24%) and 4<sup>th</sup> week of August in FH-115 (54.09%). The minimum reduction of whitefly was observed in 2<sup>nd</sup> week of July in MNH-552 (23.56%), 2<sup>nd</sup> week of August in SLH-284 (28.43%) and 1<sup>st</sup> week of July in IRFH-901 (30.06%). So the use of *C. carnea* as bio-intensive IPM program reduced the insecticides and saved foreign exchange that spent on pesticide import.

**Key Words:** Whitefly; Biological control; Bio-intensive IPM; Predator; Cotton

## INTRODUCTION

Outbreaks of cotton whitefly severely affect the economics of several cotton producing countries such as Pakistan due to direct damage by inhibition of photosynthetic activity and transmission of viral diseases to cotton that impairs the fiber quality (Henneberry *et al.*, 1999; Ahmad, 1999). Commonly three management strategies have been reported including pesticide evaluation, biological control (the evaluation of augmentative releases of commercially available natural enemies) and cultural control to suppress whiteflies (Hoddle & Robinson, 2004). Twenty three species of predators reported from cotton field including Coccinellids, Chrysopids, Lagaeids and formicids (Cheema *et al.*, 1980). *Chrysoperla carnea* is a common species with voracious polyphagous and active larvae (Kareim, 1998) has got a considerable attention as a biological control agent because of its ability to control a variety of insect pests having higher searching ability and wide adaptability in field than other predators because it attack wide host range both adults and soft bodied pests of crops (Morrison, 1985). *Chrysoperla carnea* is used in augmentative programs for sustainable crop pest suppression (Gautam & Tesfaye, 2002). The objective of the present experiment was to find out the performance of *Chrysoperla carnea* for the control of cotton whitefly to reduce the use of pesticides.

## MATERIALS AND METHODS

Six cotton cultivars, viz. MNH-552, CIM-707, SLH-


284, IRFH-901, CIM-496 and FH-115 were grown at PARS, Faisalabad in 2005 in randomized complete block design (RCBD) having two treatment groups for each variety with three replications.

Egg cards of *Chrysoperla carnea* obtained from bio-control laboratory of Department of Agri. Entomology, University of Agriculture, Faisalabad. These cards were installed in all treated replications @ 30,000 eggs/acre approximately after the interval of fifteen days during the crop season. The crop was surveyed daily for appearance of whitefly which observed on 01-07-2005. There afterwards, it was visited after an interval of a week's time till 15-09-2005 when presence of whitefly could no longer cause economic injury. Three plants selected randomly from each replication and population of whitefly counted from upper, middle and lower portion of each plant in both treated and untreated plot. The average population of whitefly per leaf was considered to be an indirect reflection of pest resistance in plants and role of bio control agent in controlling insect pests, under reference, in these studies. To see the effectiveness of bio-control agent, mean population of whitefly under study in treated (biological control) plot was compared with that of untreated (control) plot. The difference between treatment means for various parameters were subjected to Duncan's Multiple Range Test (Steel & Torrie, 1980) at 5% level of probability.


## RESULTS AND DISCUSSION

**MNH-552.** The population of whitefly was significantly


**Fig. 1. Trend of Whitefly Population in MNH-552**




**Fig. 2. Trend of Whitefly Population in CIM-707**



**Fig. 3. Trend of Whitefly Population in IRFH-901**



**Fig. 4. Trend of Whitefly Population in SLH-284**



maximum in both treated and untreated plots up to 4<sup>th</sup> week of July (Fig. 1). This population decreased gradually on the subsequent weeks of observations and the minimum population observed 1.2 and 2.8/leaf in treated and untreated

Fig. 5. Trend of Whitefly Population in CIM-496




Fig. 6. Trend of Whitefly Population in FH-115





Fig. 7. Reduction of Whitefly population due to *Chrysoperla carnea* released plots (Treated Plots) during different weeks of observation



plots, respectively. The activity of *C. carnea* found maximum in 4<sup>th</sup> week of August, resulted in maximum reduction of whitefly 57.35% and followed by 57.14, 48.68, 44.49 and 43.87% on 2<sup>nd</sup> week of September, 2<sup>nd</sup> week of August, 1<sup>st</sup> week of August and 4<sup>th</sup> week of July, respectively. Therefore, abovementioned periods favored *C. carnea* to prey maximum, whereas 2<sup>nd</sup> week of July showed minimum reduction of whitefly 23.56%, which exhibited the minimum activity of *C. carnea* for predation (Fig. 7).

**CIM-707.** The population of whitefly recorded maximum on 1<sup>st</sup> week of July both in treated (6.2/leaf) and in untreated plots (9.5/leaf), which did not differ significantly

with the whitefly population recorded up to 4<sup>th</sup> week of July in treated plot and 2<sup>nd</sup> week of July in untreated plots (Fig. 2). The minimum whitefly population was recorded to be 2.3 per leaf and 4.1 per leaf during 2<sup>nd</sup> week of September. *C. carnea* predated maximum on 2<sup>nd</sup> week of September that resulted in 45.14% reduction in whitefly population followed by 44.43, 38.57, 36.55, 36.20 and 35.31% on 1<sup>st</sup> week of September, 4<sup>th</sup> week of August, 2<sup>nd</sup> week of August, 3<sup>rd</sup> week of August and 1<sup>st</sup> week of July, respectively. The minimum activity of *C. carnea* observed on 1<sup>st</sup> week of August resulting in minimum reduction of whitefly 30.89% (Fig. 7).

**IRFH-901.** Maximum population (6.7 and 9.5 per leaf) was recorded in treated and un-treated plots, respectively on 1<sup>st</sup> week of July (Fig. 3). This population decreased gradually on the subsequent weeks both in treated and un-treated plots and reached to minimum of 2.6 and 5.1 per leaf respectively on 2<sup>nd</sup> week of September. A little variation found in population reduction of whitefly on various weeks of observation with maximum reduction in whitefly population 49.66% on 2<sup>nd</sup> week of September while minimum reduction of 30.06% on 1<sup>st</sup> week of July (Fig. 7).

**SLH-284.** A continuous and gradual decreasing trend in whitefly population per leaf observed both in treated and un-treated plots during various weeks of observation (Fig. 4) with maximum whitefly population recorded 6.3 and 8.9 per leaf in treated and un-treated plots, respectively on 1<sup>st</sup> week of July. This population was minimum on 2<sup>nd</sup> week of September 2.733 and 4.6 per leaf in treated and un-treated plots, respectively. The reduction of whitefly population recorded maximum on 1<sup>st</sup> week of September 41.98% followed by 40.58 and 38.80% on 2<sup>nd</sup> week of September and 4<sup>th</sup> week of August, respectively. The minimum reduction in whitefly population recorded 28.43% on 2<sup>nd</sup> week of August (Fig. 7).

**CIM-496.** The maximum population of whitefly recorded 6.20 and 8.90 per leaf in treated and un-treated plots respectively on 1<sup>st</sup> week of July (Fig. 5). This population decreased gradually on the subsequent weeks of observation in treated plots and un-treated plot. However, minimum population of whitefly recorded 2.4 and 4.3 per leaf in treated and un-treated plots, respectively in 2<sup>nd</sup> week of September. Maximum reduction in whitefly population observed 44.18% on 2<sup>nd</sup> week of September followed by 43.47, 40.32, 38.66 and 36.18% on 2<sup>nd</sup> week of August, 3<sup>rd</sup> week of August, 1<sup>st</sup> week of August and 4<sup>th</sup> week of July, respectively. The minimum activity of *C. carnea* observed on 1<sup>st</sup> week of July resulting in minimum population reduction 30.33% (Fig. 7).

**FH-115.** The results revealed highly significant difference among weeks both in treated and un-treated plots regarding whitefly population (Fig. 6). The maximum whitefly population recorded 6.4 per leaf on 1<sup>st</sup> week of July in treated plot and this population decreased gradually thereafter on the subsequent weeks. The minimum population observed 2.5 per leaf on 2<sup>nd</sup> week of September. The whitefly population recorded on 1<sup>st</sup> week of July was 10.3 per leaf in un-treated plots and this population increased thereafter and reached to 11.3 per leaf on 2<sup>nd</sup> week of July. The decreasing trend in population of whitefly was observed thereafter on the subsequent weeks and reached to a minimum of 4.3 per leaf on 2<sup>nd</sup> week of September. Variations were found in the data of percent reduction in whitefly population on various weeks 37.57% on 1<sup>st</sup> week of July and this reduction increased thereafter and reached up to 44.44% on 2<sup>nd</sup> week of July. The reduction in whitefly population was decreased on 3<sup>rd</sup> week of July 40.40% and again this reduction increased and reached up to 46.80% on

4<sup>th</sup> week of July. However the maximum reduction of 55.24% was observed on 3<sup>rd</sup> week of August (Fig. 7).

It was evident from the results that the effectiveness of *C. carnea* varied in different genotypes on various weeks of observation regarding whitefly population. *C. carnea* exhibited maximum effectiveness on 4<sup>th</sup> week of August for MNH-552, 1<sup>st</sup> week of September for SLH-284 and 3<sup>rd</sup> week of August for FH-115, whereas the maximum population reduction of whitefly was observed on 2<sup>nd</sup> week of September for MNH-552, CIM-707 and IRFH-901. *C. carnea*, being the generalist predator, consumes sucking pests (Chang, 1998) including 216-950 nymphs and adults of aphids and 510 nymphs of whitefly (Gautam & Tasfaye, 2002). Efficient pest management is accomplished by conservation of this predator and by managing immigrating adult population (Karahroudi & Hatami, 2003). Since half of the yield has been lost owing to recurring and imprudent appliance of chemicals so as to outcome in development resistance (Georghious & Lagunes, 1991), at that juncture, biological control is the paramount alternative, sustained by the augmentation of *C. carnea* in the containment of sucking pests particularly cotton whitefly *Bemisia tabaci* (Mohyuddin *et al.*, 1997). Consequently, it is need of the day to persuade biological control based IPM by demonstrating and recommending it on large scale.

## REFERENCES

- Ahmad, Z., 1999. Pest Problems of Cotton: A Regional Perspective. *Proceedings of ICAC-CCRI, Regional Consultation Insecticide Resistance Management in Cotton*, June 28-July 1<sup>st</sup>, Multan (Pakistan), pp: 5-20
- Chang, 1998. *Chrysoperla plurabund* larvae feed disproportionately on thrips in the field. *Canadian Entomol.*, 130: 549-50
- Cheema, M.A., N. Muzaffar and M.A. Ghani, 1980. Investigation on phenology, distribution, host range and evaluation of predators of *Pectinophora gossypiella* in Pakistan. *Pakistan Cott.*, 24:140-176.
- Gautam, R.D. and A. Tesfaye, 2002. Potential of green lacewing, *Chrysoperla carnea* (Stephens) in crop pest management. *New Agric.*, 13: 147-58
- Georghious, G.P. and T.A. Lagunes, 1991. *The Occurrence of Resistance to Pesticides in Arthropods*, p: 318. Food Agriculture Organization, United Nations, Rome
- Henneberry, T.J., N. Prabhaker, S.J. Castle and N.C. Toscano, 1999. A brief review of *Bemisia tabaci*, its occurrence and resistance to insecticides in Southern California and Arizona. In: *Proceedings of ICAC-CCRI, Regional Consultation Insecticide Resistance Management in Cotton*, June 28-July 1<sup>st</sup>, Multan (Pakistan), pp: 21-31
- Hodde, M.S. and L. Robinson, 2004. Evaluation of factors influencing augmentative releases of *Chrysoperla carnea* for control of *Scirtothrips perseae* in California avocado orchards. *Biological Control*, 31: 268-75
- Karahroudi, Z.R. and B. Hatami, 2003. Comparison of two method of releasing *C. carnea* (steph) eggs against *A. gossypii* under green house condition. *J. Sci. Tech. of Agric. Nat. Resour.*, 7: 215-25
- Kareim, A.I., 1998. Searching rate and potential of some natural enemies as bio-control agent against the *Bemisia tabaci* (Homoptera: Aleyrodidae). *J. Appl. Entomol.*, 122: 487-92
- Mohyuddin, A.I., G. Jillani, A.G. Khan, A. Hamza, I. Ahmad and Z. Mahmood, 1997. Integrated pest management of major cotton pests by conservation, redistribution and augmentation of natural enemies in Pakistan. *Pakistan J. Zool.*, 29: 293-8

Morrison, R.K., 1985. *C. carnea* In: Singh, R. and R.F. Moore (eds.), *Hand Book of Insect Rearing*, pp: 419–26. Elsevier, Amsterdam, The Netherlands

Steel, R.G.D. and J.H. Torrie, 1980. *Principles and Procedures of Statistics: A Biometrical Approach*, pp: 232–51. McGraw Hill Book Co., Inc., N.Y

**(Received 10 December 2007; Accepted 01 March 2008)**