



Full Length Article

Genetic Diversity in Three River Populations of the Giant Freshwater Prawn (*Macrobrachium rosenbergii*) in Bangladesh Assessed by Microsatellite DNA Markers

Sadequr Rahman Khan¹, Habiba Akter², Nargis Sultana², Mohd. Golam Quader Khan², Md. Abdul Wahab¹ and Md. Samsul Alam^{2*}

¹Department of Fisheries Management, Bangladesh Agricultural University, Mymensingh, Bangladesh

²Department of Fisheries Biology and Genetics, Bangladesh Agricultural University, Mymensingh-2202, Bangladesh

*For correspondence: E-mail: samsul_bau@yahoo.com

Abstract

Information on intra- and inter-population genetic diversity of cultured species may help develop efficient management and stock improvement programs. The study was undertaken to assess the genetic diversity in three wild populations namely the Paira, the Naaf and the Pashur rivers of the giant freshwater prawn (*Macrobrachium rosenbergii* de Man). Varied number of alleles, 5-14 was detected at the seven polymorphic microsatellite loci with an average of 9.833, 9.167 and 10.167 in the Paira, Naaf and Pashur populations, respectively. The average heterozygosity observed (H_o : 0.634) and expected (H_e : 0.827) were highest in the Pashur population. The H_o values of all the loci were found to be lower than the respective H_e values and thus fixation index (F_{is}) values were positive in all the populations ranging from 0.203 to 0.268, reflecting a high level of inbreeding. Ten of the 21 tests comprising seven loci in three populations for deviations from Hardy-Weinberg expectation were significant. The population differentiation (F_{ST}) value ranged from 0.012 to 0.021 indicating a low level of differentiation among the populations. Results revealed a moderate level of genetic variability though there was deficiency in observed heterozygosity in the three river populations of *M. rosenbergii*. The detection of the reducing trend in the effective population size in the rivers indicates that a pragmatic management plan is needed to establish equilibrium between the catch and the maintenance of genetic diversity of the populations of *M. rosenbergii*. © 2014 Friends Science Publishers

Keywords: *Macrobrachium rosenbergii*; Microsatellite; Heterozygosity; Genetic diversity

Introduction

The giant freshwater prawn (*Macrobrachium rosenbergii* De Man) is naturally distributed in whole of South and South-East Asia, Northern Australia and the Western Pacific Islands (New, 2002). Rearing of freshwater prawn by collecting or trapping juveniles in tidal ponds had been an age-old practice in the Indian sub-continent and Malaysia (Wickins, 1976). However, with the technologies for captive breeding and postlarvae (PLs) production established, the freshwater prawn has appeared as one of the major commercial aquaculture species in many countries such as China, India, Thailand, Vietnam and Bangladesh. Bangladesh entered into commercial prawn farming in 1990s and has acquired a position in the top five prawn producing nations of the world (FAO, 2009).

Naturally, after mating the gravid females of *M. rosenbergii* migrate towards brackish water where the eggs hatch as free-swimming larvae (Aquacop, 1977). After passing through 8 (Ling, 1969) to 11 (Uno and Soo, 1969) developmental stages, PLs move upstream to freshwater area where they live up to their adult life. Though started

with the wild-caught PL and PL collection has not been completely stopped, prawn farming in Bangladesh at present mostly depends on hatchery produced seed. However, for producing PL in the hatcheries, egg-bearing female prawns are collected from different rivers mostly from the south-west, south and south-east part of the country across the coastal districts. Therefore, the river populations are serving two important purposes: maintaining their own populations and providing mother prawn for the hatcheries. As the rivers are important reservoirs of this valuable prawn species, it is essential to conserve their genetic diversity. The genetic diversity has a positive correlation with the evolutionary potential and fitness of a population (Reed and Frankham, 2003; Vandewoestijne *et al.*, 2008). Estimation of the distribution of subpopulations based on population genetic structure may help protect weaker populations through regulating harvest based on their genetic properties (Utter, 1991). Genetic characterization also serves as a useful tool for evaluating the direction and magnitude of changes in genetic structure of species over time. Populations with a high level of genetic variation have greater prospects for

improvements in production traits such as viability, growth rate, fecundity and resistance to environmental stress and disease. Genetic variation is the basic resource that is exploited in a successful breeding program. Therefore, conservation of genetic variation is considered as an essential component of efficient fisheries management programs.

Microsatellite DNA markers are widely used for genetic characterization of wild and cultured populations for management and breeding purposes. For example, establishing pedigrees and estimating heritabilities are very effective for a breeding program that can be conducted by using microsatellite DNA markers (Norris *et al.*, 2000; Vandeputte *et al.*, 2004). In freshwater prawn and marine shrimp, microsatellite markers have been used for studying genetic diversity (Chand *et al.*, 2005; Chareontawee *et al.*, 2007; Divu *et al.*, 2008; Schneider *et al.*, 2012), for monitoring genetic variability in a breeding program (Cruz *et al.*, 2004), for assigning parentage to progeny (Jerry *et al.*, 2006) and for identifying strain as an alternative to physical tagging (Karaket *et al.*, 2011).

Detailed information on genetic structure of wild populations of *M. rosenbergii* generated by microsatellite DNA markers could play valuable roles in conserving genetic resources and managing capture fisheries as well as identifying potential source population for the development of broodstock. By applying microsatellite DNA markers, we report here for the first time the genetic population structure of three important river populations of *M. rosenbergii* from the South-West, South and South-Eastern part of Bangladesh that are used for collection of brooders and PL.

Materials and Methods

Pond and Hapa Preparation

Nine experimental ponds each with an area of 100 m² (0.01 hectare) were prepared by draining out of water and removing weeds and predators. Agricultural lime (CaCO₃) was applied at the rate of 250 kg ha⁻¹ on the bottom of the pond and the pond was filled up with underground water using a deep tube well. For promoting algal growth, the ponds were fertilized with urea, Triple Super Phosphate (TSP) and cow dung at the rates of 25 kg ha⁻¹, 25 kg ha⁻¹ and 750 kg ha⁻¹ respectively. Nine *hapas*, made of fine meshed nylon net, each having 10 m² in area (5 m × 2 m) with 1 m height were set up by using bamboo poles maintaining the top of the net 1ft above the water level.

Collection, Stocking and Management of Post-larvae (PLs) of *M. rosenbergii*

PLs were collected from three rivers namely the Pashur, the Paira and the Naaf under Bagerhat, Barguna and Cox's Bazar district, respectively (Fig. 1) and transported to the Fisheries Field Laboratory Complex, Faculty of Fisheries, Bangladesh Agricultural University, Mymensingh,

Bangladesh. After conditioning for three days, the PLs with an average weight of 0.23 g were stocked at a rate of 3 PL m⁻² (30 PL/*hapa*) with three replications. Supplementary feed having 32% protein was applied at 10% of total body weight up to 40 days and then the feeding ration was adjusted to 5% of the body weight for the next 30 days. The feed was applied twice a day, half in the morning and the other half in the afternoon.

Collection of Tissue Sample and Extraction of Genomic DNA

After 70 days of rearing, the juvenile prawns from the three *hapas* under each river source were mixed. The mean individual harvest weights (g) of the juvenile prawn were 1.51, 0.90 and 1.30 g in the Pashur, Paira and Naaf population, respectively. The pleopods of 36 randomly collected samples from each of the three river sources were clipped by using scissors and then preserved in 95% ethanol and stored at -20°C. The genomic DNA was extracted from approximately 30 mg of the tissue following standard phenol-chloroform-isoamyl alcohol extraction and ethanol precipitation method (Ayub *et al.*, 2008). The quality of the DNA samples was checked by electrophoresis on 1% agarose gel and the quantity was measured using a spectrophotometer (Biophotometer Plus, Eppendorf, Germany).

PCR Amplification

Initially, ten pairs of microsatellite primers (*Mbr1*, *Mbr2*, *Mbr3*, *Mbr4*, *Mbr5*, *Mbr7*, *Mbr8*, *Mbr9*, *Mbr10*, *Mbr11*) developed by Charoentawee *et al.* (2006) from *M. rosenbergii* were screened but finally seven pairs were used for genetic diversity assessment of three river populations of the freshwater prawn. The PCR was conducted using 50 ng template DNA, 2.0 μM of each primer, 0.25 mM of each dNTPs, 1X PCR buffer and 1 unit of Taq DNA polymerase (GENEI, India) in a total volume of 12 μL. A gradient thermocycler (Eppendorf, Germany) was used to conduct the PCR under the following temperature profiles: an initial denaturation step at 94°C followed by 35 cycles each consisting of denaturation at 94°C for 30 sec, annealing at 56°C for 30sec and extension at 72°C for 1 min. A final step of elongation at 72°C for 5 min followed the last cycle. For confirmation of PCR amplification, the products were mixed with 3 μL of loading dye (0.25% Bromophenol blue, 0.25% Xylene cyanol, 30% glycerol) and one half was run on 2% agarose gel.

Polyacrylamide Gel Electrophoresis and Silver Nitrate Staining

For separation of the microsatellite alleles, the PCR products were electrophoresed on 6% denatured polyacrylamide gel (19:1:polyacrylamide:bisacrylamide) containing 7 M urea using Sequigen GT (Bio-Rad, Hercules, USA) vertical gel electrophoresis system.



Fig. 1: Map of Bangladesh showing *M. rosenbergii* PL collection sites from three rivers

The silver nitrate staining protocol of Promega (Madison) was followed for staining the polyacrylamide gel fixed on the glass plate.

Statistical Analysis of Microsatellite Data

The length of the alleles against the molecular weight markers were estimated using the software Alpha Ease FC (Version 4.0). The genotype data on all loci of all samples of the three populations were compiled in a matrix using the Microsoft Excel program. Analysis of various population genetic parameters such as the number of alleles (N_a), effective number of alleles (N_e), frequency of alleles, observed (H_o) and expected heterozygosity (H_e), Nei's (1972) genetic distance and gene flow (N_m) between the population pairs and chi-square test for fit to Hardy-Weinberg Expectation (HWE) and Analysis of Molecular Variance (AMOVA) were performed using the software GenAlEx version 6.4 (Peakall and Smouse, 2006). The software ARLEQUIN version 3.0 (Excoffier *et al.*, 2005) was used to test the statistical significance between the F_{ST} values of different population pairs applying 1000 permutations of the genotypes. The software MEGA version 4 of Tamura *et al.* (2007) was used to build a UPGMA (unweighted pair-group method of averages) dendrogram.

Results

Allele Size and Frequency

A total of 75 alleles ranging from 220 to 338 bp were detected at the seven microsatellite loci (*Mbr-1*, *Mbr-2*, *Mbr-3*, *Mbr-4*, *Mbr-5*, *Mbr-8* and *Mbr-9*). The number of alleles varied between 5 (*Mbr-8*) and 14 (*Mbr-3*) at the seven loci with an average of 10.21 alleles per locus.

Table 1: Allele frequencies at three microsatellite loci in three populations of *Macrobrachium rosenbergii*

Locus	Allele	Paia	Naaf	Pashur	
<i>Mbr1</i>	260	0.000	0.100	0.067	
	268	0.200	0.133	0.067	
	272	0.000	0.033*	0.000	
	284	0.067	0.067	0.300	
	290	0.033	0.200	0.133	
	294	0.067	0.033	0.100	
	298	0.100	0.033	0.133	
	302	0.233	0.233	0.033	
	306	0.167	0.100	0.167	
	314	0.133	0.067	0.000	
	<i>Mbr2</i>	296	0.000	0.014	0.014
		300	0.042	0.014	0.042
		304	0.056	0.153	0.069
		308	0.111	0.056	0.208
312		0.194	0.083	0.111	
316		0.125	0.208	0.111	
320		0.250	0.222	0.139	
324		0.097	0.153	0.194	
328		0.083	0.097	0.069	
334		0.042	0.000	0.028	
338		0.000	0.000	0.014*	
<i>Mbr3</i>		234	0.083	0.097	0.111
		238	0.042	0.111	0.153
		240	0.250	0.208	0.167
	244	0.000	0.222*	0.000	
	248	0.250	0.000	0.236	
	252	0.000	0.083*	0.000	
	256	0.125	0.000	0.125	
	258	0.125	0.194	0.097	
	262	0.042	0.014	0.028	
	268	0.014	0.014	0.014	
	272	0.014	0.014	0.028	
	276	0.028	0.028	0.028	
	280	0.014	0.014	0.000	
	284	0.014	0.000	0.014	
<i>Mbr4</i>	280	0.000	0.028	0.042	
	284	0.139	0.222	0.125	
	288	0.014	0.028	0.056	
	292	0.056	0.056	0.028	
	296	0.069	0.028	0.028	
	300	0.056	0.083	0.111	
	304	0.056	0.069	0.111	
	309	0.139	0.069	0.097	
	312	0.194	0.222	0.264	
	318	0.097	0.069	0.028	
	322	0.097	0.083	0.042	
	326	0.083	0.042	0.069	
	<i>Mbr5</i>	288	0.056	0.056	0.097
		292	0.181	0.319	0.097
296		0.083	0.153	0.167	
300		0.306	0.125	0.278	
304		0.139	0.236	0.139	
308		0.069	0.028	0.056	
312		0.097	0.042	0.069	
318		0.014	0.000	0.028	
322		0.028	0.028	0.028	
328		0.028	0.014	0.042	
<i>Mbr8</i>		252	0.014	0.083	0.222
		256	0.181	0.306	0.250
		258	0.431	0.403	0.389
		262	0.278	0.153	0.111
	266	0.097	0.056	0.028	
	<i>Mbr9</i>	238	0.014	0.000	0.069
		240	0.056	0.125	0.125
		244	0.167	0.278	0.236
		248	0.306	0.347	0.306
		252	0.292	0.083	0.139
		256	0.042	0.028	0.014
		259	0.028	0.042	0.000
		262	0.028	0.056	0.014
		268	0.028	0.028	0.014
272		0.014	0.014	0.028	
276		0.014	0.000	0.028	
280		0.014	0.000	0.014	
284		0.000	0.000	0.014*	

*Indicates private alleles in the respective population

A total of five alleles were detected as private alleles: three in the Naaf population and two in the Pashur population. No private allele was detected in the Paira population (Table 1).

Intra-population Genetic Variation

A 100% of the loci were polymorphic (P_{95}) in the three studied populations. The mean number of alleles (N_a), mean effective number of alleles (N_e) and the observed (H_o) and expected (H_e) heterozygosity values of the Pashur population were higher than those of the Paira and the Naaf population (Table 2). The F_{is} (inbreeding coefficient) values were positive at all the loci in all the populations. The F_{is} value of the Pashur population was the lowest and that of the Paira population was the highest. χ^2 tests revealed deviations from the Hardy-Weinberg expectation (HWE) at varied number of loci in the three populations due to deficiencies in heterozygosity: five in the Naaf, three in the Paira and two in the Pashur population (Table 2).

Inter-population Genetic Diversity

The population differentiation value (F_{ST}) between the Paira and the Naaf population was the highest (0.021) and that between the Paira and Pashur population was the lowest (0.012) (Table 3). However, the F_{ST} values between the population-pairs were low and non-significant ($P>0.05$). Highest and lowest gene flow (N_m) values were observed between the Paira-Pashur (20.910) and the Naaf-Pashur population-pairs (11.887), respectively (Table 3). AMOVA revealed that of the total molecular variation, 6% existed among the stocks while 94% existed within stocks (data not shown).

Genetic Distance and Dendrogram

The genetic distance between the population-pairs ranged from 0.110 to 0.200. The highest and lowest genetic distances were estimated between the Pashur and Naaf population and between the Pashur and Paira population respectively (Table 3). The UPGMA dendrogram built on the basis of Nei's (1972) genetic distance included the Pashur and the Paira populations in one cluster and the Naaf population another cluster (Fig. 2).

Discussion

The abundance of alleles and heterozygosity are extensively used as tools for assessing the intra- and inter-population genetic diversity. Genetic variability is fundamental to ensuring the evolutionary potential and fitness of populations (Koljonen *et al.*, 2002; Reed and Frankham, 2003). The present study revealed a moderate level of genetic variability in terms of average number of alleles and expected heterozygosity at seven microsatellite loci in the three river populations of *M. rosenbergii*. We detected 75 alleles at seven loci (*Mbr1*, *Mbr2*, *Mbr3*, *Mbr4*, *Mbr5*, *Mbr8* and *Mbr9*) averaging 10.71 alleles per locus and their sizes

Table 2: Allelic variation and deviation from Hardy-Weinberg expectations at seven microsatellite loci in a sample of 36 *M. rosenbergii* of the three populations: N- Number of sample, N_a - number of alleles, N_e - effective number of alleles, H_o -observed heterozygosity, H_e -expected heterozygosity, F_{is} - inbreeding coefficient; H-W test- χ^2 values followed by the degrees of freedom in parenthesis

Locus	Parameters	Population		
		Paira	Naaf	Pashur
<i>Mbr2</i>	N_a	8	10	8
	N_e	6.250	6.923	5.769
	H_o	0.733	0.600	0.800
	H_e	0.840	0.856	0.827
	F_{is}	0.127	0.299	0.032
	H-W Test	35.77(28) ^{ns}	82.28(45) ^{***}	21.74 (28) ^{ns}
	Na	9	9	11
	Ne	6.612	6.276	7.261
	Ho	0.583	0.611	0.611
	He	0.849	0.841	0.862
<i>Mbr3</i>	F_{is}	0.313	0.273	0.291
	H-W Test	62.40 (36) ^{**}	57.74 (36) [*]	75.66 (55) [*]
	Na	12	11	11
	Ne	5.945	6.216	6.803
	Ho	0.611	0.556	0.583
	He	0.832	0.839	0.853
	F_{is}	0.265	0.338	0.316
	H-W Test	61.78 (66) ^{ns}	72.94 (55) ^{ns}	68.75 (55) ^{ns}
	Na	11	12	12
	Ne	8.583	7.448	7.513
<i>Mbr4</i>	Ho	0.611	0.667	0.611
	He	0.883	0.866	0.867
	F_{is}	0.308	0.230	0.295
	H-W Test	129.00 (55) ^{***}	76.96 (66) ^{ns}	133.44 (66) ^{***}
	Na	10	9	10
	Ne	5.838	4.918	6.480
	Ho	0.556	0.500	0.722
	He	0.829	0.797	0.846
	F_{is}	0.330	0.372	0.146
	H-W Test	66.36 (45) [*]	79.04 (36) ^{***}	48.74 (45) ^{ns}
<i>Mbr5</i>	Na	5	5	5
	Ne	3.281	3.461	3.620
	Ho	0.528	0.639	0.556
	He	0.695	0.711	0.724
	F_{is}	0.241	0.101	0.232
	H-W Test	15.775 (10) ^{ns}	42.540 (10) ^{***}	16.103 (10) ^{ns}
	Na	12	9	12
	Ne	4.670	4.408	5.226
	Ho	0.556	0.583	0.722
	He	0.786	0.773	0.809
<i>Mbr8</i>	F_{is}	0.293	0.246	0.107
	H-W Test	76.48 (66) ^{ns}	57.61 (36) [*]	32.69 (66) ^{ns}
	Average no. of alleles (N_a)	9.57	9.28	9.85
	Average no. of effective alleles (N_e)	5.883	5.664	6.096
	Average H_o over loci	0.574	0.592	0.634
	Average H_e over loci	0.812	0.804	0.827
	Average F_{is}	0.268	0.265	0.203
	Polymorphism (P_{95})	1.00	1.00	1.00

Key: ns=not significant, * $P<0.05$, ** $P<0.01$, *** $P<0.001$

were comparable to the findings of Chareontawee *et al.* (2006). Variations in the number of alleles have been observed in *M. rosenbergii* by different authors; e.g., 12-18 (Chand *et al.*, 2005), 5-17 (Chareontawee *et al.*, 2006),

4-20 (Chareontawee *et al.*, 2007), 3-12 (Bhassu *et al.*, 2008), 3-16 (Divu *et al.*, 2008), 10-20 (Karaket *et al.*, 2011) and 3-25 (Schneider *et al.*, 2012). The number of alleles observed in our study ranged from 5 to 14 were consistent with those of Chareontawee *et al.* (2006) and Divu *et al.* (2008) but higher than reported in a wild population in India (Schneider *et al.*, 2012). The Pashur population showed better variation than the other two populations in terms of the number of alleles and average effective number of alleles.

The average number of alleles, H_o and H_e obtained in the present study were lower than those reported in three strains of *M. rosenbergii* in Thailand (Karaket *et al.*, 2011). In all cases the H_e s were higher than the respective H_o s. H_e is considered as gene diversity as it is the expected probability that an individual will be heterozygous at a given locus. It depends on the number and relative frequencies of the alleles present in a locus (Hale *et al.*, 2012). The average observed and expected heterozygosities were found to be 0.574 to 0.634 and 0.804 to 0.827, respectively are in agreement with the range reported by other authors (Chareontawee *et al.*, 2007; Schneider *et al.*, 2012). Inbreeding causes decrease in observed heterozygosity but does not affect the allele frequencies and thus the expected heterozygosity of a population. Significant departures from HWE have been observed in all three populations- at three loci in the Paira population, at five loci in the Naaf and at two loci in the Pashur population due to reduction in heterozygosity through inbreeding. Similar to our observation, Chareontawee *et al.* (2006) also reported higher levels of H_e compared to respective H_o in a natural population of *M. rosenbergii* in Thailand. These may be attributed to the presence of null alleles in the loci examined. Deficiencies in heterozygosity relative to HWE may result from several factors such as differences in chances of mating among individuals, reduction in effective breeding number due to over exploitation (Schneider *et al.*, 2012), or selection pressure on certain locus (Ferguson, 1995; Garcia de Leon *et al.*, 1997), Wahlund effect (Hartl and Clark, 1997) and preferences of the individuals to mate with geographically nearer individuals. Among these, reduction in effective population is the most likely factor causing nonconformity to HWE in the giant prawn populations in Bangladesh. The excess of homozygosity resulted in positive F_{is} values in all populations, ranging from 0.032 to 0.372 that were higher than those reported by Schneider *et al.* (2012). F_{is} is a measure of the frequencies of heterozygotes compared to that expected when the populations are under H-W equilibrium.

The F_{ST} values between the population-pairs ranged from 0.012 to 0.021 in the present study. This level of genetic differentiation between the pairs of populations is considered as low (Balloux and Lugon-Moulin, 2002). The lowest value of F_{ST} (0.012) between the Paira and the Pashur population indicated relatively closer relation between these rivers over the Naaf population. The hierarchy of molecular

Table 3: Multilocus F_{ST} (above diagonal) and N_m (above diagonal in parentheses) and genetic distance (below diagonal) values between pairs of three stocks of *M. rosenbergii* across all loci

Population	Paira	Naaf	Pashur
Paira	***	0.021 (12.436)	0.012 (20.910)
Naaf	0.197	***	0.020 (11.887)
Pashur	0.110	0.200	***




Fig. 2: Unweighted Pair Group Method of Averages (UPGMA) dendrogram based on Nei's (1972) genetic distances between the three population pairs of *M. rosenbergii*

variance tested by AMOVA revealed that 94% of the variation existed within the population and 6% existed among the populations, which suggesting little differentiation among the studied populations. The higher level of gene flow was found between the Paira and the Pashur populations. The higher genetic distance between the Naaf and the Pashur and between the Naaf and the Paira populations may be explained by the geographical distance between the pairs as the distance of the Naaf river from the other two rivers are more than the distance between the Pashur and the Paira River (Fig. 1).

The average number of allele was lowest in the Naaf population (9.28), while average number of allele of the Pashur population (9.85) was higher with highest average effective number of alleles ($N_e = 6.096$). The effective number of alleles is much less than the actual number of alleles indicating that the frequencies are not equal for all alleles.

The genetic structure of a population is not a stable phenomenon and subjected to change over time. The magnitude and trend of the change largely depend on the intensity of natural and human interventions. The parent prawn and PL are harvested from the rivers and the sizes of the existing populations are determined by the intensity of catch and recruitment. The present study detected a reducing trend in the effective population size of *M. rosenbergii* in the rivers as reflected by deficiencies in heterozygotes and increasing inbreeding (F_{IS} values). Genetic diversity of smaller populations declines at a faster rate than that of a larger population and loss of genetic variation is considered to be the loss of fitness of a population.

In conclusion, among seven microsatellite loci Pashur population showed highest average number of alleles, and observed and expected heterozygosity. Moreover, Hardy-Weinberg expectation found fewer deviations in Pashur

population compared to the other two populations. So, genetic characterization elucidated Pashur population's PLs are superior over the Paira and the Naaf population in terms of genetic variability. These findings and catch data of last few years could help develop a pragmatic management plan to establish equilibrium between the catch and the maintenance of genetic diversity of the populations.

Acknowledgements

The research was carried out by a grant from DeLPHE (Development of Partnership in Higher Education) to MA Wahab. The equipment used in study were received from Alexander von Humboldt Foundation through an Equipment Grant to MS Alam.

References

- Aquacop, A., 1977. *Macrobrachium rosenbergii* (De man) culture in Polynesia: progress in developing a mass intensive larval rearing technique in clear water. *Proceedings of the Eighth Annual Meeting of the World Mariculture Society*, pp: 311–319. Costa Rica, 10-13 Janvier 1977. <http://archimer.ifremer.fr/doc/00000/4410/>
- Ayub, F., M.Y. Sarker and M.S. Alam, 2008. Prevalence of White Spot Syndrome Virus (WSSV) infection detected by one-step and nested PCR in selected tiger shrimp (*Penaeus monodon*) hatcheries. *Aquacult. Int.*, 16: 405–415
- Balloux, F. and N. Lugon-Moulin, 2002. The estimation of population differentiation with microsatellite markers. *Mol. Ecol.*, 11: 155–165
- Bhassu, S., L. See, M.R. Hassan, S.S. Siraj and S.G. Tan, 2008. Isolation and characterization of microsatellite loci in the Malaysian giant freshwater prawn, *Macrobrachium rosenbergii*. *Mol. Ecol. Resour.*, 8: 983–985
- Chand, V., M. De Bruyn and P.B. Mather, 2005. Microsatellite loci in the Eastern form of the giant freshwater prawn (*Macrobrachium rosenbergii*). *Mol. Ecol. Notes*, 5: 308–310
- Chareontawee, K., S. Poompuang and U. Na-Nakorn, 2006. Isolation and characterization of microsatellites in giant freshwater prawn *Macrobrachium rosenbergii*. *Mol. Ecol. Notes*, 6: 823–825
- Chareontawee, K., S. Poompuang, U. Na-Nakorn and W. Kamonrat, 2007. Genetic diversity of hatchery stocks of giant freshwater prawn (*Macrobrachium rosenbergii*) in Thailand. *Aquaculture*, 271: 121–129
- Cruz, P., A.M. Ibarra, R.H. Mejia, P.M. Gaffney and E.R. Perez, 2004. Genetic variability assessed by microsatellite in breeding program of Pacific White Shrimp (*Litopenaeus vannamei*). *Mar. Biotechnol.*, 6: 157–164
- Divu, D., I. Karunasagar and I. Karunasagar, 2008. Microsatellite DNA markers in the giant freshwater prawn, *Macrobrachium rosenbergii*: a tool for genetic analysis. *Mol. Ecol. Resour.*, 8: 1040–1042
- Excoffier L., G. Lava and S. Schneider, 2005. ARELEQUIN ver.3.0: An integrated software package for population genetics data analysis. *Evol. Bioinformatics Online*, 1: 47–50
- FAO, 2009. *Fishstat Plus (v. 2.32) issued 02.03.2009*. FAO, Rome, Italy
- Ferguson, A., 1995. Molecular genetics in fisheries: current and future prospective. In: *Molecular Genetics in Fisheries*, pp: 111–115. Carvalho, G.R. and T.J. Pitcher (eds.). Chapman and Hall, London.
- Garcia de Leon, F.J., L. Chikhi and F. Bonhomme, 1997. Microsatellite polymorphism and population subdivision in natural populations of European sea bass *Dicentrarchus labrax* (Linnaeus 1758). *Mol. Ecol.*, 6: 51–62
- Hale, M.L., T.M. Burg and T.E. Steeves, 2012. Sampling for Microsatellite-Based Population Genetic Studies: 25 to 30 Individuals per Population Is Enough to Accurately Estimate Allele Frequencies. *PLoS ONE*, 7: 1–10
- Hartl, D.L. and A.G. Clark, 1997. *Principles of Population Genetics*, 3rd edition, p: 542. Sinauer Associates, Inc, Sunderland, Massachusetts
- Jerry, D.R., B.S. Evans, M. Kenway and K. Wilson, 2006. Development of a microsatellite DNA parentage marker suite for black tiger shrimp *Penaeus monodon*. *Aquaculture*, 255: 542–547
- Karaket, T., S. Poompuang, U. Na-Nakorn, W. Kamonrat, E.M. Hallerman, 2011. DNA microsatellite-based evaluation of early growth performance among strains of freshwater prawn *Macrobrachium rosenbergii* de Man. *Aquaculture*, 311: 115–122
- Koljonen, M.L., J. Tahtinen, M. Saisa and J. Koskiniemi, 2002. Maintenance of genetic diversity of Atlantic salmon (*Salmo salar*) by captive breeding programmes and the geographic distribution of microsatellite variation. *Aquaculture*, 212: 69–92
- Ling, S.W., 1969. The general biology and development of *Macrobrachium rosenbergii* (De Man). *FAO Fish. Report*, 57: 589–606
- Nei, M., 1972. Genetic distance between populations. *Amer. Nat.*, 106: 283–292
- New, M.B., 2002. Farming freshwater prawns: a manual for the culture of the giant river prawn (*Macrobrachium rosenbergii*). FAO Fisheries Technical Paper No. 428. FAO, Rome, Italy
- Norris, A.T., D.G. Bradley and E.P. Cunningham, 2000. Parentage and relatedness determination in farmed Atlantic salmon (*Salmo salar*) using microsatellite. *Aquaculture*, 182: 73–83
- Peakall, R. and P.E. Smouse, 2006. GENEALX V6.4: Genetic Analysis in Excel. Population genetic software for teaching and research. *Mol. Ecol. Notes*, 6: 288–295
- Reed, D.H. and R. Frankham, 2003. Correlation between fitness and genetic diversity. *Conserv. Biol.*, 17: 230–237
- Schneider, K.J., J.H. Tidwell, B. Gomelsky, K.W. Pomper, G.C. Waldbieser, E. Saillant and P.B. Mather, 2012. Genetic diversity of cultured and wild populations of the giant freshwater prawn *Macrobrachium rosenbergii* (de Man, 1879) based on microsatellite analysis. *Aquac. Res.*, 44: 1425–1437
- Tamura K., J. Dudley, M. Nei and S. Kumar, 2007. MEGA4: Molecular and Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol. Biol. Evol.*, 24: 1596–1599
- Uno, Y. and K.C. Soo, 1969. Larval development of *Macrobrachium rosenbergii* (De Man reared in the laboratory. *J. Tokyo Univ. Fish.*, 55: 179–190
- Utter, F.M., 1991. Biochemical genetics and fishery management: a historical perspective. *J. Fish. Biol.*, 39: 1–20
- Vandeputte, M., M. Kocour, S. Mager, M. Dupont-Nivet, D. De Guerry, M. Rodina, D. Gela, D. Vallod, B. Chevassus and O. Linhart, 2004. Heritability estimates for growth-related traits using microsatellite parentage assignment in juvenile common carp (*Cyprinus carpio* L.). *Aquaculture*, 235: 223–236
- Vandewoestijne, S., N. Schtickzelle and M. Baguette, 2008. Positive correlation between genetic diversity and fitness in a large, well-connected metapopulation. *BMC Biol.*, 6: 46
- Wickins, J.F., 1976. Prawn biology and culture. In: *Oceanography and Marine Biology: An Annual Review*, Vol. 14, pp: 435–507. Barnes, H. (ed.). Aberdeen University Press, Aberdeen, UK

(Received 01 April 2013; Accepted 04 October 2013)